

Store at
-20°C
#69924**β-Amyloid Matched Antibody Pair**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

Species Cross Reactivity: H
UniProt ID: #P05067
Entrez-Gene Id: #351

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Isotype/Source
β-Amyloid (E3B4I) Rabbit mAb (BSA and Azide Free)	15854	100 µg	Rabbit IgG
β-Amyloid (D3D2N) Mouse mAb (BSA and Azide Free)	37372	100 µg	Mouse IgG1

Description

The β-Amyloid Matched Antibody Pair is ideal for use with immunoassay technologies and high throughput ELISA platforms requiring antibody pairs with specialized or custom antibody labeling. Labels include fluorophores, lanthanides, biotin, and beads. Platforms requiring conjugated Matched Antibody Pairs include MSD, Quanterix Simoa, Alpha Technology (AlphaScreen, AlphaLISA, LANCE, HTRF), and Luminex.

Learn how Matched Antibody Pairs move your projects forward, faster at cst-science.com/matched-antibody-pairs.

Specificity/Sensitivity

This kit detects proteins from the indicated species, as determined through in-house testing, but may also detect homologous proteins from other species.

Storage

Store at -20°C. *This product will freeze at -20°C so it is recommended to aliquot into single-use vials to avoid multiple freeze/thaw cycles.* A slight precipitate may be present and can be dissolved by gently vortexing. This will not interfere with antibody performance.

Directions for Use

Matched Antibody Pairs include capture and detection antibodies to non-overlapping epitopes. Optimal dilutions/concentrations should be determined by the end user.

Formulation

Supplied in 1X PBS (10 mM Na₂HPO₄, 3 mM KCl, 2 mM KH₂PO₄, and 140 mM NaCl (pH 7.8)). BSA and Azide Free.

Background

Amyloid β (Aβ) precursor protein (APP) is a 100-140 kDa transmembrane glycoprotein that exists as several isoforms (1). The amino acid sequence of APP contains the amyloid domain, which can be released by a two-step proteolytic cleavage (1). The extracellular deposition and accumulation of the released Aβ fragments form the main components of amyloid plaques in Alzheimer's disease (1). APP can be phosphorylated at several sites, which may affect the proteolytic processing and secretion of this protein (2-5). Phosphorylation at Thr668 (a position corresponding to the APP695 isoform) by cyclin-dependent kinase is cell-cycle dependent and peaks during G2/M phase (4). APP phosphorylated at Thr668 exists in adult rat brain and correlates with cultured neuronal differentiation (5,6).

Background References

1. Selkoe, D.J. (1996) *J Biol Chem* 271, 18295-8.
2. Caporaso, G.L. et al. (1992) *Proc Natl Acad Sci USA* 89, 3055-9.
3. Hung, A.Y. and Selkoe, D.J. (1994) *EMBO J* 13, 534-42.
4. Suzuki, T. et al. (1994) *EMBO J* 13, 1114-22.
5. Ando, K. et al. (1999) *J Neurosci* 19, 4421-7.
6. Iijima, K. et al. (2000) *J Neurochem* 75, 1085-91.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.